



UNIVERSIDADE ESTADUAL DE CAMPINAS

Instituto de Biologia

CAMILA INÉS ZORNOSA TORRES

**COMUNICAÇÃO ACÚSTICA EM ANUROS COM ESPECIAL ENFOQUE PARA:
EVOLUÇÃO DE PERDA DA VOCALIZAÇÃO DE ANÚNCIO E PRESENÇA DE
VOCALIZAÇÃO DE CORTE EM *Bokermannohyla luctuosa***

**ACOUSTIC COMMUNICATION IN ANURANS FOCUS ON: EVOLUTION OF THE
LOSS OF ADVERTISEMENT VOCALIZATION AND PRESENCE OF COURTSHIP
VOCALIZATION IN *Bokermannohyla luctuosa***

CAMPINAS

2016

CAMILA INÉS ZORNOSA TORRES

“COMUNICAÇÃO ACÚSTICA EM ANUROS COM ESPECIAL ENFOQUE PARA:
EVOLUÇÃO DE PERDA DA VOCALIZAÇÃO DE ANÚNCIO E PRESENÇA DE
VOCALIZAÇÃO DE CORTE EM *Bokermannohyla luctuosa*”

“ACOUSTIC COMMUNICATION IN ANURANS FOCUS ON: EVOLUTION OF THE
LOSS OF ADVERTISEMENT VOCALIZATION AND PRESENCE OF COURTSHIP
VOCALIZATION IN *Bokermannohyla luctuosa*”

Dissertação apresentada ao Instituto de Biologia
da Universidade Estadual de Campinas como parte
dos requisitos exigidos para a obtenção do título
de Mestre em Ecologia

Dissertation presented to the Instituto de Biologia
of the University of Campinas in partial
fulfillment of the requirements for the degree of
Master in Ecology

ORIENTADOR: Prof. Dr. Luís Felipe de Toledo Ramos Pereira

ESTE EXEMPLAR CORRESPONDE À VERSÃO FINAL DA DISSERTAÇÃO DA
ALUNA CAMILA INÉS ZORNOSA TORRES, ORIENTADA PELO PROF. LUÍS FELIPE
TOLEDO.

CAMPINAS

2016

Agência(s) de fomento e nº(s) de processo(s): CAPES; FAPESP, 2014/23388-7; CNPq, 302589/2013-9

Ficha catalográfica
Universidade Estadual de Campinas
Biblioteca do Instituto de Biologia
Mara Janaina de Oliveira - CRB 8/6972

T636c Torres, Camila Inés Zornosa, 1988-
Comunicação acústica em anuros com especial enfoque para : evolução de perda da vocalização de anúncio e presença de vocalização de corte em *Bokermannohyla luctuosa* / Camila Inés Zornosa Torres. – Campinas, SP : [s.n.], 2016.

Orientador: Luís Felipe de Toledo Ramos Pereira.
Dissertação (mestrado) – Universidade Estadual de Campinas, Instituto de Biologia.

1. Bioacústica. 2. Filogenia. 3. Anfíbio – Reprodução. 4. Comunicação animal. 5. Acasalamento de animais. I. Toledo, Luís Felipe, 1979-. II. Universidade Estadual de Campinas. Instituto de Biologia. III. Título.

Informações para Biblioteca Digital

Título em outro idioma: Acoustic communication in anurans focus on : evolution of the loss of advertisement vocalization and the presence of courtship vocalization in *Bokermannohyla luctuosa*

Palavras-chave em inglês:

Bioacoustics

Phylogeny

Amphibians –Reproduction

Animal communication

Courtship in animals

Área de concentração: Ecologia

Titulação: Mestra em Ecologia

Banca examinadora:

Luís Felipe de Toledo Ramos Pereira [Orientador]

Ivan Sergio Nunes Silva Filho

Vinícius Matheus Caldart

Data de defesa: 19-07-2016

Programa de Pós-Graduação: Ecologia

Campinas, Julho 19 de 2016

COMISSÃO EXAMINADORA

Prof. Dr. Luís Felipe de Toledo Ramos Pereira (Orientador)

Prof. Dr. Ivan Sergio Nunes Silva Filho

Dr. Vinícius Matheus Caldart

Os membros da Comissão Examinadora acima assinaram a Ata de Defesa, que se encontra no processo de vida acadêmica do aluno.

Dedico este trabalho aos meus pais, César e Claudia, e meus irmãos Lina e Alejandro, por todo o apoio durante esta caminhada longe de casa. E ao Auro por todo o amor e apoio nos piores e melhores dias.

AGRADECIMENTOS

Primeiramente gostaria de agradecer ao meu orientador o Prof. Felipe Toledo, por me receber no seu laboratório, mesmo sem ter me conhecido muito. Por sempre me apoiar e ajudar com as decisões tomadas neste projeto, por me ensinar dia a dia coisas novas e por todo o carinho com que me tratou sempre.

Gostaria também de agradecer aos membros da minha pré-banca, Dra. Cynthia Prado, Dr. Itamar Martins e Dr. Ariovaldo Giaretta, cujos comentários e sugestões ajudaram a melhorar esta dissertação para que a informação valiosa que ela contém fosse transmitida de uma forma mais clara. Também, gostaria de agradecer aos membros da banca por aceitar o meu convite e avaliar o meu trabalho.

Quero agradecer também a Dona Maria Beatriz e o Fernando por me permitir desenvolver o meu trabalho de campo na propriedade deles, á pousada Lua Cheia por me acolher sempre com muito carinho e ao Silvio pela ajuda tão importante nas saídas de campo.

Este projeto não teria sido possível sem a ajuda dos meus colegas de laboratório, Simone, Leandro, Rose, Guilherme, Sandra, Alexandre, Tami, Tsu, Carol, Carlão, Lucas e Luisa, que além de fazer comentários e sugestões para o trabalho, me ajudaram no campo ou pelo menos mostraram interesse (viram mais coletas para compartilhar) e foram um apoio nos dias de estresse contribuindo com atividades desestressantes (importantíssimas).

Também gostaria de agradecer à minha família Abadá Capoeira, já que graças a vocês consegui me relaxar, melhorar como pessoa e como capoeirista, e conhecer muito mais da cultura brasileira. Além desta grande família que está no mundo todo, agradeço aos que se converteram na minha família aqui no Brasil (Caro Valencia, Javier, Delicada, Zizi, Grazi, Pirce, Castor, Simone, Leandro, André e claro o Auro), sem vocês esses anos longe de casa não teriam sido tão prazerosos como foram, obrigada por me ensinar dia a dia o valor da amizade.

Esta vivência no Brasil e a oportunidade de estar terminando meu mestrado não teria sido possível sem o apoio da minha família, meus pais Claudia e César e os meus irmãos Lina e Alejandro, que sempre me motivaram para empreender esta aventura e me apoiaram em todo momento sem importar a distância. E aos meus amigos, Ana, Dani, Laura, Mafe, Alejandra,

Lina, Miguel, David, Natalia e Laura por compartilhar comigo este processo e a minha alegria diária.

No fim, mas não menos importante, gostaria de agradecer ao Auro, que com amor e paciência me ajudou no dia a dia, me apoio nos momentos mais difíceis, foi meu suporte nos momentos de estresse e o meu motivo de felicidade sempre, sem ele este trabalho não seria o que é neste momento, obrigada pelas ajudas com os erros do R, pelas correções no português, pelos abraços de ânimo e por todo o seu amor.

Este trabalho não poderia ter sido concluído sem o apoio financeiro da CAPES e da FASPESP (processo: #2014/23388-7)

RESUMO

A reprodução é uma das características mais conspícuas da biologia dos anfíbios, tendo uma grande diversidade em estratégias reprodutivas, modos reprodutivos e tipos de comunicação. A comunicação acústica é provavelmente a mais comum nos anuros e dentro desta, a vocalização mais estudada é o canto de anúncio. Os machos geralmente usam essa vocalização para atrair fêmeas para reprodução, as vezes utilizam o canto de corte para estimular e guiar as fêmeas ao sítio de reprodução. Apesar de estas vocalizações serem fundamentais para a maioria de anuros, algumas espécies não emitem o canto de anúncio e, portanto são consideradas mudas. Frequentemente, a ausência deste canto é atribuída ao alto ruído de fundo, sítios permanentes de reprodução, ou a falta de espécies relacionadas no mesmo habitat. Porém, estas sugestões nunca foram testadas num contexto evolutivo. Além disso, o comportamento de corte é pouco estudado nas espécies de anuros tropicais, devido à alta diversidade e à dificuldade de observar dito comportamento no campo. Assim, avaliamos se o ruído do ambiente poderia ser uma pressão seletiva das espécies ou populações de anuros mudos e descrevemos o comportamento de corte e os cantos de anúncio e corte da espécie *Bokermannohyla luctuosa*. Para a primeira parte deste estudo, revisamos as espécies mudas e mapeamos a sua distribuição na filogenia mais completa dos anuros. Estimamos o estado ancestral para o caractere mudo e analisamos a correlação entre o ruído do ambiente e a perda do canto de anúncio de duas famílias (Bufonidae e Hylidae). Em ambas as famílias, a perda do canto é dependente do ruído do ambiente. Assim, o ruído do ambiente provavelmente atua como uma pressão seletiva na perda do canto nos bufonídeos e nos hílídeos. Para a segunda parte relacionada com o comportamento reprodutivo e a comunicação de *B. luctuosa*, observamos a população de Botucatu, gravando as vocalizações com um gravador TASCAM DR680 e filmando os comportamentos usando câmeras filmadoras Sony DCR-SR47 e Sony DCR-TRV460. *Bokermannohyla luctuosa* apresenta um comportamento de corte complexo, que envolve sinais táteis e dois tipos de vocalizações (canto de anúncio e de corte), ambas as vocalizações apresentam estrutura pseudo-pulsada. O canto de corte é semelhante ao de anúncio, porém é mais longo e tem uma frequência dominante menor. A maioria das características do corte da *B. luctuosa* já foram relatadas para outras espécies da mesma família.

ABSTRACT

Reproduction is the most conspicuous characteristic in amphibian biology, with a huge diversity of reproductive strategies, modes and types of communications. Acoustic communication is probably the most common communication in anurans, and, within it, the most studied vocalization is the advertisement call. Males generally use this call to attract females for reproduction, sometimes another call (courtship call) is used to stimulate and guide females to the reproduction site. In spite of these calls being fundamental for most anurans, some species lack the advertisement call and, therefore, are considered mute. Frequently, voicelessness is attributed to high background noise, permanent breeding sites, or lack of closely related species in the same habitat. However, such suggestions have never been tested in an evolutionary context. Also, courtship behavior is poorly studied in species from the tropics, due to the high diversity and the difficulty to observe that behavior in the field. Thereby, we evaluated whether habitat noise could be a selective pressure leading to mute species or populations and describe the courtship behavior and the advertisement and courtship call of the species *Bokermannohyla luctuosa*. For the first part, we reviewed the mute species and mapped their distribution onto the most complete anuran phylogeny. We estimated ancestral states of the mute character and tested for correlation between the habitat noise and the voice loss in two families (Hylidae and Bufonidae). In both families, voice loss was dependent of habitat noise as said in a widespread assumption, environmental noise probably acted as a selective pressure on voice loss in bufonids and hylids. For the second part related with the courtship behavior of *B. luctuosa*, we observed the population from Botucatu, recording the vocalizations with a TASCAM DR680 sound recorder and filming the behaviors using a Sony DCR-SR47 and a Sony DCR-TRV460 video camera. *Bokermannohyla luctuosa* presents a complex courtship that involves tactile cues and two types of vocalizations (advertisement and courtship calls), both calls have pseudo-pulsed structure. Courtship call was similar to advertisement call but had a larger duration and a lower dominant frequency. Most of the characteristics of *B. luctuosa* courtship have also been reported for other species of the same family.

SUMÁRIO

Introdução geral	12
CAPÍTULO 1: “DOES ENVIRONMENTAL NOISE DRIVE THE LOSS OF ADVERTISEMENT CALL IN ANURANS?”	14
INTRODUCTION.....	16
METHODS	18
<i>Data acquisition and character mapping</i>	18
<i>Ancestral Character Estimation</i>	19
<i>Influence of abiotic environmental noise on voicelessness</i>	19
RESULTS	20
DISCUSSION	21
ACKNOWLEDGMENTS.....	23
FIGURES	27
CAPÍTULO 2: “COURTSHIP BEHAVIOUR OF <i>Bokermannohyla luctuosa</i> (HYLIDAE, COPHOMANTINAE) AND GEOGRAPHIC VARIATION OF ADVERTISEMENT CALL”	33
INTRODUCTION.....	36
METHODS	37
RESULTS	39
<i>Courtship and clutches</i>	39
<i>Vocalizations</i>	40
DISCUSSION	40
ACKNOWLEDGMENTS.....	42
FIGURES	46
SUPPLEMENTARY MATERIAL	50
Referências	52
ANEXOS	61

Introdução geral

A reprodução dos anfíbios é altamente variável em modos reprodutivos e compreende desde a atração dos pares até a oviposição e o cuidado parental, quando existente (Duellman & Trueb, 1996). Um dos componentes mais importantes na reprodução dos anfíbios é a comunicação. Hailman (1977) estabelece a comunicação como a transferência de informação por meio de um sinal e por um canal entre um emissor e um receptor. Esta é reconhecida pela variação no comportamento do receptor, na presença ou ausência do sinal (Hailman, 1977). Os anfíbios apresentam diferentes tipos de comunicação (acústica, visual, química, sísmica e tátil) em diversas etapas da reprodução, como a atração de parceiros e defesa de sítio de vocalização ou sítio de desova (Duellman & Trueb, 1996; Palermo-Neto & Alves, 2010).

Particularmente, para os anuros a comunicação acústica é a mais difundida e estudada. Essa comunicação está presente na vasta maioria das espécies e é frequentemente utilizada para a atração de parceiros reprodutivos, defesa territorial, defesa contra predadores e competição intrasexual (Wilczynski & Chu, 2001). Já foram reconhecidas diferentes vocalizações em anuros durante a época reprodutiva como: o canto de anúncio, principalmente emitido por machos para atrair a fêmea ou alertar outros machos da ocupação do sítio de canto (Haddad, 1995; Duellman & Trueb, 1996; Wilczynski & Chu, 2001); e o canto de corte, usado como comunicação de curta distância previa ao amplexo, para estimulação e orientação das fêmeas (Toledo *et al.* 2015).

A comunicação tátil em anuros é geralmente utilizada durante o amplexo ou a corte. Já foi relatado o uso destes sinais na identificação sexual, estimulando o amplexo de algumas espécies (*Lithobates sylvaticus*, *Hyla andersonii*, *Ascaphus truei*, *Pipa pipa*) (Duellman & Trueb, 1996). Esses sinais também foram observados envolvendo toques tanto dos machos, quanto das fêmeas durante a corte, como sinal de aceitação do amplexo (Haddad & Sawaya, 2000; Carvalho Jr. *et al.* 2006, Zina & Haddad, 2007; Lima *et al.* 2014). Esse é o caso, por exemplo, de *Aplastodiscus leucopygius*, no qual a fêmea, após de ser atraída pelo macho, toca os membros posteriores dele com as mãos ou com a cabeça, e o macho responde a esse sinal virando na direção dela e tocando-a na cabeça com a região gular ou com o focinho (Haddad *et al.* 2005).

Além de emitirem sinais unimodais, é possível também a comunicação multimodal em anuros, que se refere à produção de mais de um tipo de sinal de maneira simultânea ou

secuencial que pode possuir significado único ou variado (Narins *et al.* 2003; Preininger *et al.* 2009; Grafe *et al.* 2012; Preininger *et al.* 2012; Preininger *et al.* 2013a; Preininger *et al.* 2013b). As vezes, sinais acústicos são emitidos em consonância com visuais, como registrado em *Allobates femoralis*, *Staurois parvus*, *Sataurois latopalmatus*, *Engystomops pustulosus* (Taylor *et al.* 2008; Preininger *et al.* 2009; Grafe *et al.* 2012; Preininger *et al.* 2013a).

A ausência de vocalizações em anuros é rara e pode ser atribuída à presença de ruído de fundo intenso ou uso de locais permanentes de reprodução (Emerson & Igner, 1992). Neste caso, formas alternativas de comunicação podem ser empregadas para as espécies se comunicarem. Este é, por exemplo, o caso de *Limnonectes blythii* que se comunica visualmente para atrair a fêmea (Emerson, 1992) e *Leiopelma hamiltoni* que se comunica quimicamente para defender seu território reprodutivo (Lee & Waldman, 2002; Waldman & Bishop, 2004). No entanto, ainda não sabemos como muitas das espécies mudas se comunicam, nem se há alguma relação entre todas elas (seja ecológica ou filogenética). Neste sentido, o presente estudo realizou uma revisão bibliográfica das espécies mudas, ligando a filogenia e ecologia destas espécies e testando umas das hipóteses que explica a ausência do canto de anúncio em anuros. Além disso, realizamos a descrição do comportamento de corte e as vocalizações de anúncio e corte da espécie *Bokermannohyla luctuosa* da população de Botucatu.

**CAPÍTULO 1: “DOES ENVIRONMENTAL NOISE DRIVE THE LOSS OF
ADVERTISEMENT CALL IN ANURANS?”**



DOES ENVIRONMENTAL ABIOTIC NOISE DRIVE THE LOSS OF ADVERTISEMENT CALL IN ANURANS?

Camila Zornosa Torres*^{1,2,3}, Diogo Borges Provete⁴, Sandra Goutte^{2,3}, Luís Felipe Toledo²

¹Programa de Pós-Graduação em Ecologia, Instituto de Biologia, Universidade Estadual de Campinas, C.P. 6109, 13083-970, Campinas, São Paulo, Brazil.

²Laboratório Multiusuário de Bioacústica (LMBio), Departamento de Biologia Animal, Instituto de Biologia, Universidade Estadual de Campinas, Campinas, São Paulo, Brazil.

³Laboratório de História Natural de Anfíbios Brasileiros (LaHNAB), Departamento de Biologia Animal, Instituto de Biologia, Universidade Estadual de Campinas, Campinas, São Paulo, Brazil.

⁴CNPq Post-doctoral fellow, Department of Biological and Environmental Sciences, University of Gothenburg, Göteborg, Sweden.

*Corresponding author: cami_zt@hotmail.com

Abstract

Acoustic communication is probably the most common communication in anurans, and, within it, the most studied vocalization is the advertisement call. Mainly males use this call to attract conspecific females for reproduction. In spite of this call being fundamental for most anurans, some species lack the advertisement call and, therefore, are considered mute. Frequently, voicelessness is attributed to high background noise, permanent breeding sites, or lack of closely related species in the same habitat. However, such suggestions have never been tested in an evolutionary context. We evaluated whether habitat noise could be a selective pressure leading to mute species or populations. We reviewed the mute species and mapped their distribution onto the most complete anuran phylogeny available to date. We estimated ancestral states of the mute character. We listed 40 mute species belonging to 7 families, and suggested that the mute trait is the ancestral state in the phylogeny we use. We further investigated two families, Hylidae and Bufonidae, which have the highest number of mute species represented in the phylogenetic tree, and tested the relation between voice loss and ambient noise level. The rate of evolution of the advertisement call was dependent of habitat noise, as a widespread assumption, environmental noise probably acts as a selective pressure on voice loss in bufonids and hylids. Based on our results, we indicate the need of additional natural history studies, focusing on alternative communication modes among mute species in order to provide new insights about possible selective forces that are taking place in this case.

Keywords: Amphibian communication, ancestral reconstruction, Pagel's correlation, phylogeny, voiceless frogs evolution, background noise.

INTRODUCTION

Animal communication is central in evolutionary biology and of fundamental importance in species natural history, as it mediates reproductive, defense, territorial, aggressive, and social interactions (Noble 1998; Narins 2001; Fitch 2006; Toledo and Haddad 2009; Toledo et al. 2011). Communication signals are under constant selection for maximizing the information transmitted, providing unambiguous cues and adapting to various

habits in form of different modes of communications, all these aspects let it remain in the evolutionary history of the species (Hauser 1996).

In anurans there are at least five channels of communication: acoustic, visual, chemical, tactile, and seismic, in which acoustic is by far the most common (Vitt and Caldwell 2009). This communication mode is present at different phases of the reproduction, such as in mate attraction and defense of calling or spawning sites (Wilczynski and Chu 2001; Palermo-Neto and Alves 2010). Consequently, the acoustic channel has been recognized as the most important communication mode in anurans (Wells 2007; Dorcas et al. 2010). Vocalizations have a tremendous importance in anuran communication because they transmit relevant information, such as individual's location, size, reproductive status, aggressive motivation, or can mediate predator-prey interactions (Davis 1987; Bee and Gerhardt 2001; Ryan 2001; Simmons 2004; Wells 2007; Toledo and Haddad 2009; Toledo et al. 2015).

Fourteen different types of calls, subdivided into three main categories, have been described for anurans (Toledo et al. 2015; Köhler et al. submitted). The most common vocalization in anurans is the advertisement call, produced principally by males in the reproductive season, but it can also be produced by females (Duellman and Trueb 1994; Haddad 1995; Wilczynski and Chu 2001; Toledo et al. 2015; Preininger et al. in press). It is highly variable and can be a simple note or a series of identical notes, it can be produced in the frequency of audible sound, infrasound or ultrasound (Toledo et al. 2015). Advertisement calls can also combine those frequencies, as registered for the call of *Odorrana tormota*, which combines ultrasound and audible frequencies (Feng et al. 2006). Males produce advertisement calls to attract conspecific females or males to join the chorus; it allows them to distribute themselves within a calling site and for the female to localize them individually while it also transmits information about male size that has an important role in sexual selection and in physical combats (Toledo et al. 2015).

Given the importance of advertisement calls, it is surprising to find that some species lack such vocalizations, referred to as mute or voiceless frogs (Emerson and Inger 1992; Haddad and Giaretta 1999; Rödel et al. 2003; Wells 2007; Toledo et al. 2015). Mute species are expected to use alternative channels of communication in place of advertisement calls (Emerson 1992; Lee and Waldman 2002; Waldman and Bishop 2004). This prediction was corroborated in the case of males of *Limnodynastes blythii* (Dicroglossidae), which uses visual signaling to attract females (Emerson 1992), and males of *Leiopelma hamiltoni* (Leiopelmatidae), which communicates chemically to defend its reproductive territory (Lee and Waldman 2002; Waldman and Bishop 2004).

The absence of advertisement calls has been attributed to the presence of intense background noise, the use of permanent breeding sites, or the presence of closely related species in the same habitat (Emerson and Inger 1992). Presence of low frequency background noise could mask anurans vocalizations, which would need to use more energy to being heard. However, the hypotheses linking voicelessness and breeding environment, to our knowledge, has not been tested yet. In this study, we investigated whether species living in noisy environments have a higher propensity to lack advertisement call. To test this hypothesis we did a correlation between both characters and searched for the ancestral character state of the advertisement call.

METHODS

Data acquisition and character mapping

We defined as mute or voiceless, anurans that do not emit advertisement calls; the presence of other call types was not considered. We searched for mute anuran species in the available literature, searching the online databases and university libraries using the keywords: voiceless frog, mute frog, reproductive biology, communication, reproduction, Amphibia, ecology, and specific epithets of species suggested to be mute. Additionally, some information was gathered from personal communications. Information about habits and habitat of each species were collected from the IUCN redlist website (IUCN 2015). Taxonomy follows Frost (2016).

The presence of advertisement call was coded as a binary character (0: absent, 1: present) for each species, species having mute and calling populations were consider as calling species. After compilation of the data, we mapped advertisement call absence and their abundance on the most recent and comprehensive global amphibian phylogeny (Pyron 2014). This phylogeny includes representatives of more than 90 % of the currently recognized genera and over 3,300 amphibian species (Pyron 2014), we considered mute species, and those not mute belonging to the Bufonidae and Hylidae. To prune the species we used the function *prune.sample* of the package “picante” from the R platform (Kembel et al. 2014).

As some reports of absence of advertisement calls could result from a lack of natural history knowledge, we considered mute only those species that have been reported to be voiceless in more than one reference, or those that have been reported voiceless for a long time (more than five years). If these were not the case (e.g., just one recent reference indicates

that the species is mute), we considered the species as probably mute. We also categorized the families in ranges (0%, 0.1-5%, 5.1-50%, >50%) considering the number of mute species and the total number of species of the family.

Environmental noise was coded as binary (0: not noisy and 1: noisy). We considered noisy habitats those with waterfalls and rapid streams; all other cases (such as terrestrial, lentic environments, and slow rivulets) were considered not noisy habitats.

Ancestral Character Estimation

Based on the phylogenetic hypothesis published by Pyron (2014), we estimated ancestral presence of advertisement call in two different ways: one in separate phylogenies for bufonids and hylids, without considering mute species of other families; and the other in a unique phylogeny that considers bufonids, hylids, and mute species of these and other five families. We used the function *ace* from the R package “ape”, with the symmetrical model (SYM) (Paradis et al. 2015). The ancestral states reconstruction of discrete characters considered the maximum likelihood estimation, using the probabilities of each character states at each node of the tree, assuming that all states of the character are possible at each node (even if some are more likely or more parsimonious than others) and that each character evolves at a constant rate over the tree (Pagel 1994; Cunningham et al. 1998). The probabilities were determined by the distribution of the character state in the tip species, by the rate of character evolution and by the inter-nodal branch lengths (Cunningham et al. 1998).

Influence of abiotic environmental noise on voicelessness

In order to test whether the loss of advertisement call was related to the environmental noise, we did a Pagel’s correlation test (Pagel 1994), using information of families Bufonidae and Hylidae, and complete information of all the mute species. We used a total of 495 species pruned from the phylogeny published by Pyron (2014). The Pagel’s correlation test uses a continuous-time Markov model to characterize evolutionary changes along each of the branches of a phylogeny (Pagel 1994). This method tests the hypothesis of correlation between two binary characters by comparing the fit of two models to the observed data set.

The first model treats the two characters as evolving independently, the goodness of fit of this model is then compared to a more complex model in which the characters evolve in a correlated fashion (dependent model) (Pagel 1994). The dependent model is justified as a representation of the data if it fits the data significantly better than the independent model; their significance is assessed by means of a likelihood ratio statistic (Pagel 1994).

Thereby, we built an independent and dependent model of correlation and compared their goodness of fit to our data. We selected the best fitting model using the Akaike's Information Criterion (AIC), and the log-likelihood statistic. This analysis was implemented in the R platform using the "phytools" package and the function *fitPagel*, we also add the argument *dep.var* = "x" to fit the direction of our hypothesis (advertisement call rate of evolution depends on the environmental noise rate of evolution) (Pagel 1994; Revell 2012).

RESULTS

We listed 40 mute anuran species, assigned to 7 families, dispersed over the order Anura (Fig. 1) and globally distributed (Table 1). Leiopelmatidae is the only family with all species mute; three families have between 5.1 and 50 % of mute species, and three of the seven families have from 0.1 to 5 % of mute species. A clade with three sister genera containing mute species (*Incilius*, *Anaxyrus*, and *Rhinella*) was highlighted in the family Bufonidae (Fig. 2). Several mute species were recognized in the family Hylidae, including a genus, *Megastomatohyla*, containing only mute species (Fig. 3). Families Bufonidae and Hylidae were selected to more detailed analysis because of the high number of mute species and its representativity in the tree.

The advertisement call loss occurred four times within the Bufonidae and three times within the Hylidae, when using different phylogenies (Fig. 4), being the presence of advertisement call the ancestral state. When using the unique phylogeny of both families and including mute species of other families, the ancestral character state was the absence of the advertisement call (Fig. 5), after that the advertisement call loss eleven times within the phylogeny. The Pagel's correlation test showed that both models are different (likelihood-ratio = 4.04; $P = 0.04$). The best fitting model for our data set was the one that shows that the rate of evolution of the advertisement call depends on the rate of evolution of the environmental noise (Independent model: AIC = 544.09, Log-likelihood = -270.05; Dependent model: AIC = 542.05, Log-likelihood = -268.03).

DISCUSSION

For the first time we provided, based on a comprehensive analysis, support to the previous hypothesis that the lack of advertisement calls is related to the presence of abiotic background noise (Emerson and Inger 1992; Wells 2007). Responses to environmental noise as a selective pressure varies among anuran clades, as some lineages or clades evolved to use alternative signals or voicelessness (Rödel et al. 2003; Grafe et al. 2012; Stratenberger et al. 2013). For example, *Leiopelma* species use visual and chemical communication (Green 1988; Lee and Waldman 2002; Waldman and Bishop 2004; Bell 2010) and *Acanthixalus* spp. produce volatile substances as an alternative communication (Rödel et al. 2003; Stratenberger et al. 2013). Therefore, anurans showed to be variable in relation to the different outcomes selected under the influence of background noise.

Furthermore, some studies suggest that environmental characteristics can influence the evolution of long-range acoustic signals and in some cases can even modulate species vocalizations (Bosch and De la Riva 2004). Different vocal characteristics are linked with environmental noise. For example, some species that reproduce in streams have advertisement calls with high dominant frequency, suggesting that advertisement calls have evolved towards higher frequencies, thereby avoiding spectral overlap with background noise (Goutte et al. 2013; Röhr et al. 2015). These frogs may also use visual signaling and in some cases ear tuning acting as a frequency band filter (Goutte et al. 2013; Röhr et al. 2015; de Sá et al. 2016).

The absence of advertisement call in frogs has been considered as an scarce evolutionary event (Wells 2007), as few species were reported as mute. Nonetheless, we showed that even the number of mute species is low in proportion to the amphibian species of the world (Frost 2016), it is considerably high considering that acoustic communication is the principal communication mode in anurans (Wells 2007; Toledo et al. 2015). It is possible that we over-estimated the number of mute species, as a consequence of insufficient natural history data. On the other hand, we could be also under-estimating this number, when assuming that several species produce advertisement calls (when there is no studies that suggest the contrary). We suggest that voiceless ancestral condition can varies depending on the method used, and we propose that voiceless is an ancestral condition, as Leiopelmatidae (a family sister to all other families in the Anura clade) (Frost et al. 2006; Pyron and Wiens 2011; Pyron 2014), Gymnophiona and Caudata (other Orders of Amphibia, being caecilinas

sister to Anura: Fig. 1), lacks advertisement call (Wells 2007), agreeing to what was previously proposed (Stephenson and Stephenson 1957).

Otherwise, the loss of advertisement call in anurans evolved many times independently, as also observed for Bornean voiceless frogs that lost the advertisement call twice (Emerson and Berrigan 1993). Such high frequency of losses may imply that the advertisement call, although a very important type of communication in anurans (Wells 2007; Toledo et al. 2015), could be easily lost during evolutionary history. An intermediate state could be the loss of the advertisement call primarily in the reproductive communication, being overcome by a different communication modality. For example, visual communication plays a major role in some cases, such as in the genera *Brachycephalus* (Pombal Jr. et al. 1994; Araújo et al. 2012).

At last, additional studies focused on the reproductive biology and communication of voiceless species are necessary to assure their proper classification as mute or not. Advertisement calls and calling behavior are highly variable and species may be misclassified as mute if they produce very rare or low intensity calls (see Blair and Pettus 1954; Wells 2007; Wells 1977; Toledo et al. 2015). Some species have been incorrectly reported as mute for a long time, such as *Incilius alvarius* (Bufonidae), *Mantidactylus guttulatus* (Mantellidae), and *Insuetophrynus acarpicus* (Rhinodermatidae) (Blair and Pettus 1954; Diaz et al. 1983; Sullivan and Malmos 1994; Vences et al. 2004). Others have mute and calling populations as *Bufo bufo* (Bufonidae), *Anaxyrus boreas* (Bufonidae), *Limnonectes kuhlii* (Dicroglossidae), and *Limnonectes blythii* (Dicroglossidae) (Black and Brunson 1971; Höglund and Robertson 1988; Emerson 1992; Matsui 1995; Orlov 1997; Emerson and Ward 1998; Tsuji and Lue 1998; Wells 2007). These last cases should be target of subsequent studies, as they may provide comparative scenario (between mute and vocal populations), helping to understand better loss of call evolution. Besides this, it is noteworthy that some mute species can also emit other types of calls, such as release and distress calls (e.g., Schuierer 1962; Black and Brunson 1971; Bell 1978; Penna and Veloso 1987; Green 1988; Höglund and Robertson 1988; Jacobson and Vanderberg 1991; Toledo and Haddad 2009; Grant and Bolívar-G 2014). Therefore, the terminology mute should be avoided or clearly defined when necessary. We also suggest studies that involve anatomical evaluation of mute species.

ACKNOWLEDGMENTS

We thank Alex Pyron for providing the most recent phylogeny, Kentwood D Wells, Peter M Narins, Walter Hödl, Célio FB Haddad, Julián Faivovich, Rafael Márquez, and Mark-Oliver Rödel, for providing unpublished information about mute frogs; We also thank financial support from the Brazilian Federal Agency for Support and Evaluation of Graduate Education (CAPES), São Paulo Research Foundation (FAPESP grant #2014/23388-7), and National Counsel of Technological and Scientific Development (CNPq grant #302589/2013-9).

Table 1. The 40 mute or probably mute (indicated by an asterisk) anuran species of the world, countries where they naturally occur, breeding habitat, and references indicating that they are mute.

Species	Distribution	Breeding habitat	Reference
Alsodidae			
<i>Alsodes montanus</i> *	Chile	Streams	Penna and Veloso 1987
<i>Alsodes tumultuosus</i> *	Chile	Streams	Penna et al. 1983
<i>Alsodes valdiviensis</i> *	Chile	Streams	Formas et al. 2002
Bufonidae			
<i>Incilius periglenes</i> *	Costa Rica	Pools	Jacobson and Vanderberg 1991
<i>Incilius holdridgei</i>	Costa Rica	Streams	Novak and Robinson 1975
<i>Incilius peripatetes</i> *	Panama	Unknown	Novak and Robinson 1975
<i>Incilius epioticus</i> *	Costa Rica and Panama	Unknown	Novak and Robinson 1975
<i>Incilius fastidiosus</i>	Costa Rica and Panama	Streams	Novak and Robinson 1975
<i>Anaxyrus exsul</i>	United States of America	Pools and Streams	Schuieler 1962
<i>Anaxyrus nelsoni</i>	United States of America	Pools	Wells 2007
<i>Rhinella paraguas</i> *	Colombia	Unknown	Grant and Bolivar 2014
<i>Rhinella acrolopha</i> *	Colombia	Unknown	Novak and Robinson 1975
<i>Rhinella spinulosa</i> *	Argentina, Chile, Bolivia and Peru	Lakes, streams and pools	Penna and Veloso 1981
Hylidae			
<i>Bokermannohyla izecksohni</i>	Brazil	Streams	Jim and Caramaschi 1979
<i>Plectrohyla siopela</i>	Mexico	Streams	Wells 2007
<i>Plectrohyla</i>	Mexico	Streams	Canesco-Márquez et al.

ameibothalame 2002

Charadrahyla altipotens Mexico Streams Duellman 1970

Megastomatohyla mixomaculata Mexico Streams Wells 2007

Megastomatohyla pellita Mexico Streams Wells 2007

Megastomatohyla mixe Mexico Streams Wells 2007

Megastomatohyla nubicola Mexico Streams Wells 2007

Hylodidae

Hylodes vanzolinii Brazil Streams L. F. Toledo (unpubl. data)

Megaelosia apuana Brazil Streams Pombal et al. 2003

Megaelosia bocainensis Brazil Streams Giaretta et al. 1993

Megaelosia boticariana Brazil Streams Giaretta & Aguiar, 1998

Megaelosia goeldii Brazil Streams Giaretta et al. 1993

Megaelosia jordanensis Brazil Streams Heyer, 1983

Megaelosia lutzae Brazil Streams Giaretta et al. 1993

Megaelosia massarti Brazil Stream Giaretta et al. 1993

Hyperoliidae

Acanthixalus sonjae Ghana Tree cavities Rödel et al. 2003
Stranberger et al. 2013

Acanthixalus spinosus Cameroon, Congo, Nigeria, Gabon, Democratic Republic of the Congo Tree cavities Rödel et al. 2003
Stranberger et al. 2013

Leiopelmatidae

<i>Leiopelma archeyi</i>	New Zealand	Shallow depressions beneath logs	Bell 2010
<i>Leiopelma hamiltoni</i>	New Zealand	Unknown	Bell 2010
<i>Leiopelma hochstetteri</i>	New Zealand	Land, shallow water	Bell 2010
<i>Leiopelma pakeka</i>	New Zealand	Moist depressions under logs, rocks or vegetation	Bell 2010
Telmatobiidae			
<i>Telmatobius halli</i> *	Chile	Unknown	Penna and Veloso 1987
<i>Telmatobius marmoratus</i> *	Peru, Bolivia, Chile, Argentina	Streams, waterfalls	Penna and Veloso 1987
<i>Telmatobius pefauri</i>	Chile	Streams	Penna and Veloso 1987
<i>Telmatobius peruvianus</i> *	Chile and Peru	Streams	Penna and Veloso 1987
<i>Telmatobius zapahuirensis</i> *	Chile	Streams	Penna and Veloso 1987

FIGURES

Fig. 1 Distribution of mute anuran families (colored branches) in the anuran phylogeny, with caecilians and salamanders as outgroups (based on Pyron 2014). In parenthesis number of mute species and total number of species of each family.

Fig. 2 Distribution of mute character (colored branches) in Bufonidae. In parenthesis number of mute species and total number of species of each genus.

Fig. 3 Distribution of mute character (colored branches) in Hylidae. In parenthesis number of mute species and total number of species of each genus.

Fig. 4 Ancestral character estimation of advertisement call presence in bufonids (A) and (B) hylids in separate phylogenies without mute species of other families. Red arrows indicate branches where the mute character appears in the phylogeny.

Fig. 5 Ancestral character estimation of advertisement call presence in bufonids, hylids and mute species belonging to other families in a unique phylogeny. Black portions in the pie plot represent the probability of lacks advertisement call. Red portions in the pie plot represents the probability of have advertisement call.

Fig. 1




Fig. 2




Fig. 3






Fig. 4

Fig. 5



CAPÍTULO 2: “COURTSHIP BEHAVIOUR OF *Bokermannohyla luctuosa* (HYLIDAE, COPHOMANTINAE) AND GEOGRAPHIC VARIATION OF ADVERTISEMENT CALL”

**COURTSHIP BEHAVIOUR OF *Bokermannohyla luctuosa* (HYLIDAE,
COPHOMANTINAE) AND GEOGRAPHIC VARIATION OF ADVERTISEMENT
CALL**

Camila Zornosa Torres*^{1,2,3}, Luís Felipe Toledo³

¹Programa de Pós-Graduação em Ecologia, Instituto de Biologia, Universidade Estadual de Campinas, C.P. 6109, 13083-970, Campinas, São Paulo, Brazil.

²Laboratório Multiusuário de Bioacústica (LMBio), Departamento de Biologia Animal, Instituto de Biologia, Universidade Estadual de Campinas, Campinas, São Paulo, Brazil.

³Laboratório de História Natural de Anfíbios Brasileiros (LaHNAB), Departamento de Biologia Animal, Instituto de Biologia, Universidade Estadual de Campinas, Campinas, São Paulo, Brazil.

*Corresponding author: cami_zt@hotmail.com

Abstract

Reproduction is the most conspicuous characteristic in amphibian biology, because there is a great diversity of reproductive modes. In order to reproduce, male anurans generally display acoustic communication and complex courtship behaviors. Hylids have one of the most variable reproductive repertoires, however we lack information for many species, especially in regard to courtship and acoustic communication. Therefore, we studied *Bokermannohyla luctuosa*, a species with the reproductive behavior poorly documented. Hence, we described for the first time its complex courtship, involving tactile communication (touches between the pair) and two types of vocalizations: advertisement and courtship calls. These calls presented pseudo-pulsed structure. The courtship call was similar to advertisement call but had a larger duration, lower dominant frequency and lower relative peak power. Most of the characteristics of *B. luctuosa* courtship have also been reported for other species of the subfamily Cophomantinae, such as *Bokermannohyla nanuzae*, *Aplastodiscus leucopygius*, *A. arildae*, *A. perviridis*, and *Hypsiboas rosenbergi*.

Key words: Advertisement call, axillary amplexus, reproduction, breeding.

INTRODUCTION

Courtship is fundamental for many amphibian species, contribute their reproductive fitness (Duellman & Trueb, 1996). During courtship, amphibians use different types of signals (acoustic, visual, tactile and chemical) to attract mates. Among these, the most commonly reported are chemical (mainly in salamanders), acoustic and tactile (mainly in anurans) (Wells, 1977). Besides this, secondary structures are also important in some species, as enlargement of the cloacal glands in male salamanders or presence of nuptial excrescences in male anurans (Duellman & Trueb, 1996).

However, courtship is just one aspect of reproduction. Traditionally anuran reproductive strategies are categorized into 39 reproductive modes, that includes simple modes as eggs laid in ponds where tadpoles will develop, and more complex modes with eggs swallowed by the mother, completing the subsequent development in its stomach (Haddad & Prado, 2005). Additionally, eggs are deposited in different types of nests, besides gelatinous egg masses, such as foam or bubble nests, on the water, ground or above trees (Haddad & Prado, 2005). Furthermore, anurans vary in their development, from indirect (with tadpoles or larval stage), to direct as in the terrestrial eggs of brachycephaloids (Haddad & Prado, 2005; Wells, 2007; Haddad *et al.* 2013). In spite of such high diversity of breeding modes, many, if not most of the anurans, and especially in the Tropical region, lack information about their reproductive biology. Studies are even scarcer if the considered aspect is courtship. Therefore, such known diversity may be underestimated, highlighting the need of studying reproductive behavior in natural conditions.

Among anurans, the genus *Bokermannohyla* (Hylidae) is interesting to conduct studies of reproductive biology. As breeding biology is still poorly studied for the genus *Bokermannohyla*, the information about their courtship is even more scarce in the literature, this information is known only for 3 of the 32 species of the genus: *B. ibitiguara*, *B. nanuzae*, and *B. alvarengai* (Nali & Prado, 2012; Lima *et al.* 2014; Centeno *et al.* 2015). Species of the *Bokermannohyla circumdata* group breeds in streams, where males generally emit advertisement calls perched on the marginal vegetation or on the ground near the stream (Caramaschi & Feio, 1990; Carvalho *et al.* 2012). Courtship description is only available for *B. nanuzae*, a species that uses visual (face wiping), tactile (alternating touches in different parts of the body), and acoustic signaling (Lima *et al.* 2014).

Composition of the advertisement call in this group varies from 1 to 30 notes, lasting for up to 7.6 seconds (Napoli & Caramaschi, 2004; Carvalho *et al.* 2012; Gaiga *et al.* 2013; Lima *et al.* 2014). Courtship calls, present in at least *B. nanuzae*, differ from the advertisement calls by having higher rate of emission (from 21 to 44 notes/minute) and longer note duration (from 0.34 to 0.61 s) (Lima *et al.* 2014). Some congeneric species are commonly found in bromeliads (Napoli & Pimenta, 2003; Napoli & Caramaschi, 2004; Napoli & Juncá, 2006; Napoli & Pimenta, 2009; Carvalho *et al.* 2012). However, in most of the cases these females lay their eggs in nearby streams or ponds (Carvalho *et al.* 2012; Haddad *et al.* 2013). Besides that, data available on the reproduction of *Bokermannohyla* spp. focus on its acoustical communication during the breeding season (Carvalho *et al.* 2012; Gaiga *et al.* 2013).

For *B. luctuosa* the advertisement call has been described as composed (with two notes), with frequencies varying from 0.3 to 2.0 kHz, emitted from the ground or perched on vegetation (Pombal & Haddad, 1993; Carvalho *et al.* 2012; Gaiga *et al.* 2013). Call descriptions are available for populations of Serra do Japi (Jundiaí) and Atibaia, both localities in the state of São Paulo, Brazil. However, no information is available about its courtship behavior. Therefore, based on field observations in Botucatu (São Paulo, Brazil) we describe its courtship, including egg laying site and male-female acoustic communications before and during the amplexus. Additionally we compare the advertisement call of these three populations (Jundiaí, Atibaia and Botucatu), as a taxonomic confirmation method, because *B. luctuosa* has not been reported to Botucatu.

METHODS

We conducted field observations at “Sítio Santo Antônio da Cascatinha” located in the district of Rubião Junior, municipality of Botucatu, state of São Paulo, southeastern Brazil (22°53'42.9" S, 48°29'28" W). The study site is included in the Atlantic forest biome, has an elevation of 850 meters, and is consider a semideciduous forest fragment with native plants surrounded by an agriculture matrix (Figure 1A). Inside this fragment there is a stream with approximately 5 to 60 cm of depth, 30 to 100 cm wide, with crystalline water and sandy bottom (Figures 1B-C).

We studied the population of *Bokermannohyla luctuosa* of this fragment between November 2014 and March 2015, during its reproductive season (Pombal Jr. & Haddad, 1993) totaling 37 nights. Specimens were found by *Ad libitum* active search followed by focal animal observations from 19:00 to 3:00 (Altmann, 1974). We filmed the specimens using a Sony DCR-SR47 and a Sony DCR-TRV460 video camera, using a red light lantern or the night vision mode of the cameras. We recorded the vocalizations with a TASCAM DR680 sound recorder coupled to a Sennheiser unidirectional microphone, at a sample frequency of 48 kHz and a sample size of 24 bits.

Prior to analyses, frequencies over 3 kHz were filtered out. Calls were normalized (removing DC offset, centering on 0.0 vertically, and to the maximum amplitude of -1.0 dB) in Audacity 2.1.0. Treated files were saved as 16 bits WAV files. We then analyzed these recordings in Raven Pro 1.4 (Charif *et al.* 2010). The configurations adopted were: 70% brightness, 80% contrast, and a Fast Fourier Transform length (FFT) of 1200.

To describe the calls we used the term pseudo-pulses, here defined as a very short sound with a low frequency resolution and imperfect pulsation (Robilliard *et al.* 2006; Araújo, 2011; see also Fig. 3B). Spectral parameters were measured in the spectrogram and temporal parameters were measured in the oscillogram.

The peak of dominant frequency was obtained with the Peak Frequency function. We used the Frequency 5% and Frequency 95% functions to obtain the minimum and maximum frequencies respectively. The Frequency 5% function corresponds to the minimum frequency cutting the 5% of the energy of the call below it and the Frequency 95% corresponds to the maximum frequency cutting the 5% of the energy of the call above it (Charif *et al.* 2010). Functions Frequency 5% and Frequency 95% were used to estimate the bandwidth that concentrates 90% (BW 90%) of the energy of the call.

The relative peak power was measured in Raven only for note A in both advertisement and courtship calls. Values were corrected using relative differences in the same recording. I.e., we compared the peak power of the advertisement and courtship calls of the same recordings, subtracting from both measurements the minimum peak power measured. The same was done with three different recordings. Therefore, these values were not absolute, but fit for comparisons between different call types and recordings.

Temporal parameters as note duration, call duration and interval between notes were measured using the function Delta time. We also measure call rate manually, by counting the number of calls in one minute for each record. We did a two sample *t*-test to compare temporal parameters and the relative peak power between advertisement and courtship calls of

B. luctuosa. Recordings were deposited at Fonoteca Neotropical Jacques Viellard, Unicamp, Campinas, Brazil (FNJV 32378–86).

RESULTS

Courtship and clutches

We observed the courtship of one couple of *Bokermannohyla luctuosa*. The couple was found the dawn of March 9th of 2016, in a pool inside a shallow depression of approximate 20 cm of diameter in the margin of the stream. The observation started at 3:00 am and the courtship finished at 4:15 am. We observed the male calling in this depression for almost 13 minutes before the female approached it. At that point the male was calling oriented to the opposite direction of the female. When the female approached the male, it increased its calling rate from 19 calls/minute to 44 calls/minute. Then, the male (Figure 2A) turned toward the female and approached her. After that, the male placed his hand on female's head for 12 seconds. During that period, the female remained motionless and the male reduced the calling rate from 52 calls/minute (before placing his hand on her head) to 7 calls/minute (just after touching the female). After about five minutes the female moved closer to the male, who jumped onto her back, entering in axillary amplexus. This first amplexus lasted 57 seconds. Three seconds after releasing the female, the male amplexed the female again and emitted a short amplexant call (not considered for analysis). After 58 seconds of this second axillary amplexus the male released the female again, remained 12 seconds at her side, and amplexed her for the third time, which lasted for 21 seconds. During each of these three consecutive amplexus the female laid eggs. After this, the male pushed the eggs to the extreme of the depression and then amplexed the female for a fourth time. This action was repeated three times, lasting 50 seconds and completing six amplexus cycles. After the sixth amplexus, the female hid in the depression's border and male stayed moving the eggs with its hands for about two minutes.

We also observed three clutches in shallow depressions with leaf litter inside them. These depressions were close (5 cm) or far (from 1 to 3 m) from the stream. One observed clutch had eggs of different stages of development (Figure 2B).

Vocalizations

We recorded two distinct vocalizations: the advertisement call, emitted by males while attracting females from long distance to their position, and the courtship call, emitted by males in presence of the female. Their (advertisement) calling activity started on January at 23:00h and lasts during the remaining field period ($n = 176$ calls). All males were calling inside ground depressions in the border of the stream or in depressions under logs. We recorded nine calling males (FNJV 32377–86), but only four males of those were analyzed (FNJV 32380–81, 32384–85) due to the quality of the recordings.

Both the advertisement and the courtship calls presented pseudo-pulsed structure (Figure 3B) and were composed by two notes (Figures 3A and C) emitted in different combinations (Table 1). In both calls, note A was longer and had a higher peak of dominant frequency than note B (Table 2). Occasionally, note A when emitted in the advertisement call had a short portion at the beginning of the note separate from a longer second part (Figure 3A).

We analyzed 347 notes of advertisement call from four different males (mean = 86.75 notes/male). Advertisement calls were emitted in a mean rate of 23.5 ± 10.62 calls/minute. This call lasted on average 1.11 ± 0.62 seconds (Table 3), with an average peak of dominant frequency of 537.85 ± 276.09 Hz. We also analyzed 278 notes of courtship calls from three different males (mean = 92.6 notes/male), emitted in a mean rate of 47 ± 9.90 calls/minute. Courtship calls were longer than advertisement calls, emitted with a higher rate and with a lower relative peak power than the advertisement call (Table 3).

DISCUSSION

Acoustic signals and tactile stimuli are an important part of *Bokermannohyla* spp. courtship, in some cases the cues can be harder as in *B. ibitiguara* (where male hit the female with his arm) or a simple touch as in *B. nanuzae* (Nali & Prado, 2012; Lima *et al.* 2014). In this study we saw tactile stimuli similar to that reported for *B. nanuzae*, *Aplastodiscus leucopygius* and *A. arildae* (all included in the Cophomantinae subfamily *sensu* Duellman *et al.* 2016), in which male touches the female with his hands in different parts of female's body (head and dorsum) (Haddad & Sawaya, 2000; Carvalho Jr. *et al.* 2006, Zina & Haddad, 2007;

Lima *et al.* 2014). Such stimuli are commonly observed in species in which males construct a nest prior to female attraction (Haddad & Sawaya, 2000; Zina & Haddad, 2007). Therefore, it is possible that the depression where the couple was observed could have been previously constructed by the male of *B. luctuosa*.

From our observations we suggest that the advertisement call of *B. luctuosa* was produced to attract the female and the courtship call could stimulate the female to accept males' amplexus and also stimulate its egg laying (Toledo *et al.* 2015). Although the advertisement call structure in the *B. circumdata* group showed to be highly variable, for example, the number of notes varies from one to 30, calls lasts from 0.1 up to 7.6 seconds, and peak of dominant frequency varies from 0.3 to 2.7 kHz (Gaiga *et al.* 2013), we found similarities. In agreement to our results, the advertisement call of species of the *Bokermannohyla circumdata* group is generally composed by pulsed notes as those described for *B. ibitiguara*, *B. nanuzae* and *B. sazimai* (Eterovick & Brandão, 2001). Therefore, we suggest that this character could be shared to all of its

Geographic variations of calls were reported for *B. nanuzae*, *B. circumdata*, and *B. sazimai* (Carvalho *et al.* 2012; Gaiga *et al.* 2013). Advertisement call of *B. luctuosa* was described from the type locality (Serra do Japi, Jundiaí, São Paulo, Brazil: Pombal & Haddad, 1993) and Atibaia (São Paulo, Brazil: Carvalho *et al.* 2012). The advertisement call of population from Atibaia had two different notes emitted consecutively (Carvalho *et al.* 2012), similar to what we described from Botucatu. The mean duration of the call in these three populations (Serra do Japi, Atibaia and Botucatu) is variable (Table S1). The advertisement call of population from Serra do Japi has only one note, and is shorter than the other two populations (Pombal & Haddad, 1993). Note B is longer (0.44 ± 0.046) than note A (0.057 ± 0.02) in individuals from Atibaia (Carvalho *et al.* 2012), contrary to what we observed in Botucatu (note A: 0.77 ± 0.18 s; note B: 0.19 ± 0.05 s). Also, calls of individuals from Atibaia are higher-pitched (0.49 to 2.0 kHz) (Carvalho *et al.* 2012) than calls of the other two populations, the lowest-pitched call is from individuals from Serra do Japi (0.3 to 1.8 kHz) (Pombal & Haddad, 1993). The highest peak of dominant frequency in the population of Botucatu is from note A (1.8 kHz) and in the population from Atibaia is from note B (2.0 kHz). Therefore, we suggest that those notes could be the same as also observed in the spectrogram (Carvalho *et al.* 2012) (Table S1).

Courtship call of *B. luctuosa* was never described in previous studies, probably due to the low probability of recording such event in field activities. Many species of the Cophomantinae subfamily emit courtship calls, as *B. nanuzae*, *A. leucopygius*, *A. arildae*, *A.*

perviridis, and *Hypsiboas rosenbergi* (Kluge, 1981; Haddad & Sawaya, 2000; Haddad *et al.* 2005; Carvalho Jr. *et al.* 2006; Lima *et al.* 2014). Some species have more than one courtship call and this call may be variable and it is emitted when the female approaches the male or while guiding the female to the spawning site (Toledo *et al.* 2015).

The courtship call usually presents similar structure to the advertisement call, being the rate of emission and power the main differences. Courtship calls from *H. rosenbergi*, *A. perviridis* and *B. nanuzae* have a lower rate of emission when compared to their advertisement calls (Kluge, 1981; Haddad *et al.* 2005; Lima *et al.* 2014). On the other hand, *A. leucopygius*, *A. arildae* (Haddad & Sawaya, 2000; Carvalho Jr. *et al.* 2006) and *B. luctuosa* (present study) have a higher rate of emission in their courtship call compared with its advertisement call. Peak of dominant frequency in courtship calls is usually the same as those of advertisement calls (*B. nanuzae*, *A. leucopygius*, *H. rosenbergi*) (Kluge, 1981; Haddad & Sawaya, 2000; Lima *et al.* 2014). However, a lower peak of dominant frequency in the courtship call has been reported for *A. perviridis* (Haddad *et al.* 2005) and presently for *B. luctuosa*. We showed that the relative peak power of the courtship call is lower than the advertisement call, whilst other studies did not analyzed this parameter.

Even few is known about species of the *B. circumdata* group, some of them have been reported to use depressions or burrows near the stream as spawning sites (Nali & Prado, 2012; Lima *et al.* 2014; Centeno *et al.* 2015). However, the use of the same site for different clutches has been reported only for *B. nanuzae* (Lima *et al.* 2014) and *B. luctuosa* (Pombal & Haddad, 1993). Such particular characteristic might indicate that males of these species use the same site to attract different females. More detailed studies about reproductive behavior are needed for species of *Bokermannohyla* from different groups.

ACKNOWLEDGMENTS

We thank S. C. de Almeida, L. F. Moreno, A. N. Pimentel Jr., N. M. Novoa, V. Sacardi, S. Goutte, L. Ribeiro and C. H. L. Nunes de Almeida for helping during fieldwork; Maria Beatriz and Ferdando for letting us work in their property. Financial support was provided by the Brazilian Federal Agency for Support and Evaluation of Graduate Education (CAPES), São Paulo Research Foundation (FAPESP grant #2014/23388–7), and National Counsel of Technological and Scientific Development (CNPq grant #302589/2013–9).

Table 1. Variable composition of the advertisement call of *Bokermannohyla luctuosa* with two notes of unknown function. Values in parenthesis correspond to the percentage of frequency of the call composition in 176 calls.

Calls composition	Frequency in 176 calls
A	92 (52.3 %)
AB	27 (15.3 %)
AABB	13 (7.4 %)
ABB	12 (6.8 %)
AA	8 (4.5 %)
AAB	8 (4.5 %)
AAABB	3 (1.7 %)
AABBB	2 (1.1 %)
BAB	2 (1.1 %)
BBAA	2 (1.1 %)
AAA	1 (0.6 %)
AAABAB	1 (0.6 %)
AABAB	1 (0.6 %)
AABABBB	1 (0.6 %)
AABBAB	1 (0.6 %)
ABAAB	1 (0.6 %)
BB	1 (0.6 %)

Table 2. Acoustic parameters for the notes of the advertisement and courtship calls of males of *Bokermannohyla luctuosa*. Values presented as Mean \pm SD (range).

	Note A (N=346)		Note B (N=232)	
	Advertisement call	Courtship call	Advertisement call	Courtship call
Note duration (s)	0.73 \pm 0.18 (0.19 – 1.53)	0.84 \pm 0.17 (0.33 – 1.48)	0.19 \pm 0.05 (0.07 – 0.41)	0.19 \pm 0.04 (0.08–0.34)
Freq 5% (Hz)	359.55 \pm 21.22 (257.80 – 539.10)	412.01 \pm 96.45 (70.30 – 609.40)	304.68 \pm 75.12 (46.90 – 375.00)	327.95 \pm 37.79 (93.80 – 468.80)
Peak Freq (Hz)	653.85 \pm 433.26 (375.00 – 1757.80)	748.65 \pm 484.61 (281.20 – 1804.70)	376.95 \pm 77.44 (70.30 – 679.70)	495.11 \pm 252.24 (328.10 – 1710.90)
Freq 95% (Hz)	1863.85 \pm 341.49 (820.30 – 2671.90)	2027.18 \pm 166.66 (1429.70 – 2367.20)	1996.09 \pm 656.37 (773.40 – 2906.20)	2199.85 \pm 425.12 (1148.10 – 2976.60)
BW 90% (Hz)	1504.32 \pm 337.01 (468.80 – 2343.80)	1615.17 \pm 197.59 (1078.10 – 2015.60)	1691.41 \pm 660.21 (445.30 – 2695.30)	1871.90 \pm 425.59 (820.30 – 2718.80)

Table 3. Temporal and spectral parameters for advertisement and courtship calls of *Bokermannohyla luctuosa* and results of individual *t*-tests. Values presented as Mean \pm SD (range; sample size). See methods for the explanation of the relative peak power.

	Advertisement call (N = 4 males)	Courtship call (N = 3 males)	<i>t</i>-test
Call duration (s)	1.11 \pm 0.62 (0.52 – 3.78; 176)	16.24 \pm 15.51 (3.87 – 50.49; 10)	$t = 13.36$; df = 184; $P < 0.0001$
Intervals between notes (s)	1.72 \pm 2.43 (0.01 – 14.91; 313)	0.08 \pm 0.03 (0.02 – 0.25; 266)	$t = 9.29$; df = 577; $P < 0.0001$
Call rate (calls/minute)	23.5 \pm 10.62 (6 – 40; 14)	47 \pm 9.90 (36 – 60; 10)	$t = 5.49$; df = 22; $P < 0.0001$
Relative peak power (dB)	10.37 \pm 3.5 (4 – 16; 30)	2.49 \pm 2.80 (0 – 11.2; 30)	$t = 9.61$; df = 58; $P < 0.0001$

FIGURES

Figure 1. Study site, indicating the studied forest fragment (A) and the stream inside this fragment (B-C), at the district of Rubião Junior, municipality of Botucatu, state of São Paulo, southern Brazil.

Figure 2. Male of *Bokermannohyla luctuosa* (A), and an egg mass with eggs in different developmental stages (B).

Figure 3. Spectrogram and oscillogram of the advertisement call (FNJV 32381) of *Bokermannohyla luctuosa*, municipality of Botucatu, São Paulo, Brazil (A). Structure of pseudo-pulses in advertisement and courtship calls of *B. luctuosa* (B). Spectrogram and oscillogram of the courtship call (FNJV 32381) of *Bokermannohyla luctuosa*, municipality of Botucatu, São Paulo, Brazil (C).

Figure 1.



Figure 2.





Figure 3.



SUPPLEMENTARY MATERIAL

Table S1. Temporal and spectral parameters for advertisement call of three populations of *Bokermannohyla luctuosa*. Duration values presented as mean \pm SD (range), dominant frequency values presented as range.

Population	Note duration (s)		Dominant frequency (Hz)	Publication
	Note A	Note B		
Serra do Japi	0.06		305 – 1734	Pombal & Haddad 1993
Atibaia	0.057 ± 0.02	0.44 ± 0.046	490 – 2000	Carvalho <i>et al.</i> 2012
Botucatu	0.77 ± 0.18 (0.19 – 1.53)	0.19 ± 0.05 (0.07 – 0.41)	300 – 1800	Present study

Considerações finais

Este estudo proporcionou informações sobre as espécies de anuros mudos, encontrando 40 espécies reportadas como mudas, e a relação entre a falta da vocalização de anúncio com o ruído do ambiente. Também encontramos que os resultados da reconstrução do estado ancestral podem variar dependendo o método usado. Assim, detectamos que o estado ancestral ao testar as famílias separadamente foi a presença de canto de anúncio, contrário ao resultado obtido testando as duas famílias na mesma filogenia e inserindo todas as espécies mudas.

Além disso, este estudo forneceu conhecimentos sobre o comportamento reprodutivo de *Bokermannohyla luctuosa*, descrevendo o canto de anúncio e o canto de corte. Também descrevemos o comportamento de corte que envolve sinais acústicos e táteis, similarmente como em outras espécies da mesma subfamília. Com os nossos resultados mostramos que há uma variação geográfica no canto desta espécie, novamente como relatado para outras espécies do gênero.

Baseado nos nossos resultados, destacamos a importância de realizar futuras análises com as outras hipóteses sugeridas para a perda da vocalização de anúncio, assim como adicionar medições de ruído nos habitats das espécies mudas. Embora evidente, também vale ressaltar que a realização de estudos de história natural para as espécies de anuros para as quais ainda não temos informação sobre a presença ou ausência de vocalização é fundamental.

Referências

Introdução geral

- Carvalho Jr., R.R; Galindo, C.A.B; Nascimento, L.B. 2006. Notes on the courtship behavior of *Aplastodiscus arildae* (Cruz & Peixoto, 1985) at an urban forest fragment in southeastern Brazil (Amphibia, Anura, Hylidae). *Arquivos do Museu Nacional, Rio de Janeiro*. 64(3): 247-254.
- Duellman, W.E; Trueb, L. 1996. *Biology of Amphibians*. The Johns Hopkins University Press. Baltimore and London. 670 p.
- Emerson, S.B. 1992. Courtship and nest-building behavior of a Bornean frog, *Rana blythi*. *Copeia* 1992(4):1123-1127.
- Emerson, S.B; Igner, R.F. 1992. The comparative ecology of voiced and voiceless Bornean Frogs. *Journal of Herpetology* 26(4):482-490.
- Grafe, T.U; Preininger, D; Sztatecsny, M; Kasah, R; Dehling, J.M; Proksch, S; Hödl, W. 2012. Multimodal communication in a noisy environment: A case study of the Bornean frog *Staurois parvus*. *PLoS ONE* 7(5)
- Haddad, C.F.B. 1995. Comunicação em anuros (Amphibia). *Anais de Etologia*. 13:116-132.
- Haddad, C.F.B; Faivovich, J; Garcia, P.C.A. 2005. The specialized reproductive mode of the treefrog *Aplastodiscus perviridis* (Anura: Hylidae). *Amphibia-Reptilia* 26:87-92.
- Haddad, C.F.B & Sawaya, R. 2000. Reproductive modes of Atlantic Forest Hylids frogs: A general overview and the description of a new mode. *BIOTROPICA* 32(4): 862-871.
- Hailman, J.P. 1977. *Optical signals: Animal communication and light*. Bloomington: Indiana University Press. 362 p.
- Lee, J.S.F & Waldman, B. 2002. Communication by fecal Chemosignals in an Archaic Frog, *Leiopelma hamiltoni*. *Copeia* 3:679-686.
- Lima, N.G.S; Gontijo, A.S.B; Eterovick, P.C. 2014. Breeding behavior of *Bokermannohyla nanuzae* (Anura: Hylidae) at an Atlantic Forest site in southeastern Brazil. *Journal of Natural History* 48(23-24): 1439-1452.
- Narins, P.M; Hödl, W; Grabul, D.S. 2003. Bimodal signal requisite for agonistic behavior in a dart-poison frog, *Epipedobates femoralis*. *PNAS* 100(2):577-580
- Palermo-Neto, J; Alves, G.J. 2010. A comunicação dos animais. *Revista CFMV* 16(49):24-34.

- Preininger, D; Boeckle, M; Freudmann, A; Stranberger, I; Sztatecsny, M; Hödl, W. 2012. Multimodal signaling in the small torrent frog (*Micrixalus saxicola*) in a complex acoustic environment. *Behavioral Ecology and Sociobiology*.
- Preininger, D; Boeckle, M; Sztatecsny, M; Hödl, W. 2013a. Divergent receiver responses to components of multimodal signals in two foot-flagging frog species. *PLoS ONE* 8(1):e55367.
- Preininger, D; Boeckle, M; Hödl, W. 2009. Communication in noisy environments II: Visual signaling behavior of male foot-flagging frogs *Staurois latopalmatus*. *Herpetologica* 65(2):166-173.
- Preininger, D; Stiegler, M.J; Gururaja, K.V; Vijayakumar, S.P; Torsekar, V.R; Sztatecsny; Hödl, W. 2013b. Getting a kick out of it: multimodal signaling during male-male encounters in the foot-flagging frog *Micrixalus* aff. *Saxicola* from the Western Ghats of India. *Current Science* 105(12):1735-1740.
- Taylor, R.C; Klein, B.A; Stein, J; Ryan, M.J. 2008. Faux frogs: multimodal signaling and the value of robotics in animal behavior. *Animal Behaviour* 76(2):1089-1097.
- Toledo, L.F; Martins, I.A; Bruschi, D.P; Passos, M.A; Alexandre, C; Haddad, C.F.B. 2015. The anuran calling repertoire in the light of social context. *Acta Ethol.* 18:87–99.
- Waldman, B. & Bishop, P. J. 2004. Chemical communication in an archaic anuran amphibian. *Behavioral Ecology*: 15(1):88-93.
- Wilczynski, W. & Chu, J. Em Ryan, M. J. 2001. Anuran Communication. Smithsonian Institution Press. Washington and London. 252 p.
- Zina, J. & Haddad, C.F.B. 2007. Courtship behavior of two treefrog species, *Aplastodiscus arildae* and *A. leucopygius* (Anura: Hylidae), from the Atlantic Forest, southeastern Brazil. *Herpetological Review* 38(3): 282-285.

Capítulo 1

- Araújo CB, Guerra TJ, Amatuzzi MCO, Campos LA (2012) Advertisement and territorial calls of *Brachycephalus pitanga* (Anura: Brachycephalidae). *Zootaxa* 3302:66–67.
- Bee MA, Gerhardt HC (2001) Neighbour-stranger discrimination by territorial male bullfrogs (*Rana catesbeiana*). *An. Behav.* 62:1129–1140.
- Bell BD (1978) Observations on the ecology and reproduction of the New Zealand Leiopelmid frogs. *Herpetol.* 34(4):340–354.

- Bell B (2010) The threatened Leiopelmatid frogs of New Zealand: Natural history integrates with conservation. *Herpetol. Cons. Biol.* 5(3):515–528.
- Black JH, Brunson RB (1971) Breeding behavior of the boreal toad, *Bufo boreas boreas* (Baird and Girard), in western Montana. *Great. Basin. Nat.* 31(2):109–113.
- Blair WF, Pettus DI (1954) The mating call and its significance in the Colorado River toad (*Bufo alvarious* Girard) *Tex. J. Sci.* 6:72–77.
- Bosch J, De la Riva I (2004) Are frog calls modulated by the environment? An analysis with anuran species from Bolivia. *Can. J. Zool.* 82(6):880–888.
- Canesco-Márquez L, Mendelson JR, Gutiérrez-Mayén G (2002) A New Species of *Hyla* (Anura: Hylidae) from the Mixteca Alta, Oaxaca, Mexico. *Herpetol.* 58(2):260–269.
- Cunningham CW, Omland KE, Oakley TH (1998) Reconstructing ancestral character states: a critical reappraisal. *Trends. Ecol. Evol.* 13:361–366.
- Davis MS (1987) Acoustically mediated neighbor recognition in the North American bullfrog, *Rana catesbeiana*. *Behav. Ecol. Sociobiol.* 21:185–190.
- De Sá FP, Zina J, Haddad CFB (2016) Sophisticated communication in the Brazilian Torrent Frog *Hylodes japi*. *Plos One* 11(1):e0145444. doi:10.1371/journal.pone.0145444
- Diaz N, Valencia J, Sallaberry M (1983) Life history and phylogenetic relationships of *Insuetophrynus acarpicus* (Anura: Leptodactylidae). *Copeia* (1):30–37.
- Dorcas ME, Price SJ, Walls SC et al. (2010) Auditory monitoring of anuran populations. In: Dood CK Jr (Ed.) *Amphibian Ecology and Conservation: A handbook techniques*. Oxford: Oxford University Press. New York.
- Duellman WE (1970) The hylid frogs of Middle America. *Monogr. Mus. Nat. Hist., Univ. Kans.* Volume 2. 1:1–753.
- Duellman WE, Trueb L (1994) *Biology of Amphibians*. The Johns Hopkins University Press. Baltimore and London.
- Emerson SB (1992) Courtship and nest-building behavior of a Bornean frog, *Rana blythi*. *Copeia* 1992(4):1123–1127.
- Emerson SB, Berrigan D (1993) Systematics of Southeast Asian Ranids: multiple origins of voicelessness in the subgenus *Limnonectes*. *Herpetol.* 49(1):22–31.
- Emerson SB, Inger RF (1992) The comparative ecology of voiced and voiceless Bornean Frogs. *J. Herpetol.* 26(4):482–490.
- Emerson SB, Ward R (1998) Male secondary sexual characteristics, sexual selection and molecular divergence in fanged ranid frogs of Southeast Asia. *Zool. J. Linn. Soc.* 122:537–553.

- Feng AS, Narins PM, Xu CH et al (2006) Ultrasonic communication in frogs. *Nature* 440:333–336.
- Fitch T (2006) Production of vocalizations in mammals. Elsevier. University of St Andrews, UK
- Formas JR, Cuevas CC, Brieva LM (2002) A new species of *Alsodes* (Anura: Leptodactylidae) from Cerro Mirador, Cordillera Pelada, southern Chile. *Proc. Biol. Soc. Wash.* 115(4):708–719.
- Frost DR (2016) Amphibian Species of the World: an Online Reference. American Museum of Natural History, New York, USA. Available from <http://research.amnh.org/vz/herpetology/amphibia/> (accessed February 2016)
- Frost DR, Grant T, Faivovich J et al. (2006) The amphibian tree of life. *Bull. Amer. Mus. Nat. Hist.* 297:1–370.
- Giaretta AA, Bokermann WCA, Haddad CFB (1993) A review of the genus *Megaelosia* (Anura: Leptodactylidae) with a description of a new species. *J. Herpetol.* 27(3):276–285.
- Giaretta AA, Aguiar O Jr (1998) A new species of *Megaelosia* from the Mantiqueira range, Southeastern Brazil. *J. Herpetol.* 32(1):80–83.
- Goutte S, Dubois A, Legendre F (2013) The Importance of Ambient Sound Level to Characterise Anuran Habitat. *Plos One* 8(10):e78020. doi:10.1371/journal.pone.0078020
- Grafe TU, Preininger D, Szatcscny M et al. (2012) Multimodal Communication in a Noisy Environment: A case Study of the Bornean Rock Frog *Staurois parvus*. *Plos One* 7(5):e37965. doi:10.1371/journal.pone.0037965
- Grant T, Bolívar-G W (2014). A new species of semiarboreal toad with a salamander-like ear (Anura: Bufonidae: *Rhinella*). *Herpetol.* 70(2):198–210.
- Green DM (1988) Antipredator behavior and skin glands in the New Zealand native frogs, genus *Leiopelma*. *N. Z. J. Zool.* 15(1):39–45.
- Haddad CFB (1995) Comunicação em anuros (Amphibia). *An. Etol.* 13:116–132.
- Haddad CFB, Giaretta AA (1999) Visual and acoustic communication in the Brazilian torrent frog, *Hylodes asper* (Anura: Leptodactylidae). *Herpetol.* 55(3):324–333.
- Hauser MD (1996) The evolution of communication. The MIT Press. London, England.
- Höglund J, Robertson JGM (1988) Chorusing behavior, a density-dependent alternative mating strategy in male common toads (*Bufo bufo*). *Ethol.* 79:324–332.
- Jacobson SK, Vandenberg JJ (1991) Reproductive ecology of the endangered golden toad (*Bufo periglenes*). *J. Herpetol.* 25(3):321–327.

- Jim J, Caramaschii U (1979) Uma nova espécie de *Hyla* da região de Botucatu, São Paulo, Brasil (Amphibia, Anura). Rev. Bras. Biol. 39(3):717–719.
- Kembel SW, Cowan MR, Helmus WK et al. (2010). Picante: R tools for integrating phylogenies and ecology. Bioinformatics 26:1463–1464.
- Köhler J, Jansen M, Rodríguez A et al. The use of bioacoustics in anuran taxonomy: theory, terminology, methods and recommendations for best practice. Zootaxa (submitted)
- Lee JSF, Waldman B (2002) Communication by fecal chemosignals in an Archaic Frog, *Leiopelma hamiltoni*. Copeia 3:679–686.
- Matsui M (1995) Call produced by a “voiceless” frog, *Rana blythi* Boulenger 1920, from Peninsular Malaysia (Amphibia Anura). Trop. Zool. 8:325–331.
- Narins PM (2001) Vibration Communication in Vertebrates. In: Ecology of sensing. Department of Physiological Science, UCLA, Los Angeles, California, USA, pp 128–148.
- Noble J (1998) The evolution of animal communication systems: questions of function examined through simulation. University of Southampton, University of Sussex. PhD Thesis. 194 pp.
- Novak RM, Robinson D (1975) Observations on the reproduction and ecology of the tropical montane toad, *Bufo holdridgei* Taylor in Costa Rica. Rev. Biol. Trop. 23(2):213–237.
- Orlov N (1997) Breeding behavior and nest construction in a Vietnam frog related to *Rana bythi*. Copeia 1997(2):464–465.
- Pagel M (1994) Detecting correlated evolution on phylogenies: a general method for the comparative analysis of discrete characters. Proc. R. Soc. Lond. B 255:37–45.
- Palermo-Neto J, Alves GJ (2010) A comunicação dos animais. Rev. CFMV 16(49):24–34.
- Paradis E, Blomberg S, Bolker B, et al. (2015) Analyses of phylogenetics and evolution. Package ‘ape’ CRAN. Available from: <http://ape-package.ird.fr/> (accessed January 2016).
- Penna M, Contreras S, Veloso A (1983) Acoustical repertoires and morphological differences in the ear of two *Alsodes* species (Amphibia: Leptodactylidae). Can. J. Zool. 61:2369–2376.
- Penna M, Veloso AM (1981) Acoustical signals related to reproduction in the *spinulosus* species group *Bufo* (Amphibia, Bufonidae). Can. J. Zool. 59:54–60.
- Penna M, Veloso A (1987) Vocalization by Andean frogs of the genus *Telmatobius* (Leptodactylidae). Herpetol. 43(2):208–216.
- Preininger D, Handschuh S, Boeckle M et al. Comparison of female and male vocalisation and larynx morphology in the size dimorphic foot-flagging frog species *Staurois guttatus*. Herpetol. J. (in press).

- Pombal Jr. JP, Prado GM, Canedo C (2003) A new species of giant torrent frog, genus *Megaelosia*, from the Atlantic rain forest of Espírito Santo, Brazil (Amphibia: Leptodactylidae). *J. Herpetol.* 37(3):453–466.
- Pombal Jr. JP, Sazima I, Haddad CFB (1994) Breeding Behavior of the Pumpkin Toadlet, *Brachycephalus ephippium* (Brachycephalidae). *J. Herpetol.* 28(4):516–519.
- Pyron RA (2014) Biogeographic analysis reveals ancient continental vicariance and recent oceanic dispersal in amphibians. *Syst. Biol.* 63(5):779–797.
- Pyron RA, Wiens JJ (2011) A large-scale phylogeny of Amphibia including over 2800 species, and a revised classification of extant frogs, salamanders and caecilians. *Mol. Phylogenet. Evol.* 61:543–583.
- Revell LJ (2015) Phylogenetic tools for comparative biology (and other things). Package ‘phytools’ CRAN. Available from: <http://www.phytools.org> (accessed January 2016).
- Rödel MO, Kosuch J, Veith M et al. (2003) First record of the genus *Acanthixalus* from the upper Guinean Rain Forest, West Africa, with the description of a new species. *J. Herpetol.* 37(1):43–52.
- Röhr DL, Brant-Paterno G, Camurugi F et al. (2015) Background noise as a selective pressure: stream-breeding anurans call at higher frequencies. *Org. Divers. Evol.* doi:10.1007/s13127-015-0256-0
- Ryan MJ (2001) Anuran communication. Smithsonian Institution Press, Washington and London.
- Schneider FW (1962) Remarks upon the natural history of *Bufo exsul* Myers, the endemic toad of Deep Springs Valley, Inyo County, California. *Herpetol.* 17(4):260–266.
- Simmons AM (2004) Call recognition in the bullfrog, *Rana catesbeiana*: generalization along the duration continuum. *J. Acoust. Soc. Am.* 115(3):1345–1355.
- Stephenson EM, Stephenson NG (1957) Field Observations on the New Zealand Frog *Liopelma* Fitzinger. *T. Roy. Soc. NZ* 84(4):867–882.
- Stranberger I, Poth D, Saradhi-Peram P et al (2013) Take time to smell the frogs: vocal sac glands of reed frogs (Anura: Hyperoliidae) contain species-specific chemical cocktails. *Bio. J. Linnean Soc.* 110(4):828–838.
- Sullivan BK, Malmos KB (1994) Call variation in the Colorado River Toad (*Bufo alvarius*): behavioral and phylogenetic implications. *Herpetol.* 50(2):146–156.
- The IUCN Red List of Threatened Species. (2015) Version 2015.2. www.iucnredlist.org. (accessed March 2015)

- Toledo LF, Haddad CFB (2009) Defensive vocalizations of Neotropical anurans. *South Am. J. Herpetol.* 4(1):25–42.
- Toledo LF, Martins IA, Bruschi DP et al. (2015) The anuran calling repertoire in the light of social context. *Acta Ethol.* 18:87–99.
- Toledo LF, Sazima I, Haddad CFB (2011) Behavioural defences of anurans: an overview. *Ethol. Ecol. Evol.* 23(1):1–25.
- Tsuji H, Lue KY (1998) Temporal aspects of the amplexus and oviposition behavior of the fanged frog *Rana kyllii* from Taiwan. *Copeia* 1998(3):769–773.
- Vences M, Andreone F, Glaw F (2004) Voice of a giant: bioacoustic data for *Mantidactylus guttulatus* (Amphibia: Mantellidae). *Amphibia-Reptilia* 25:112–115.
- Vitt LJ, Caldwell JP (2009) *Herpetology*. 3rd Edition. Academic Press Elsevier. China.
- Waldman B, Bishop PJ (2004) Chemical communication in an archaic anuran amphibian. *Behav. Ecol.* 15(1):88–93.
- Wells KD (1977) The social behaviour of anuran amphibians. *Anim. Behav.* 25:666–693.
- Wells KD (2007) *The Ecology and Behavior of amphibians*. The University of Chicago Press. Chicago and London.
- Wilczynski W, Chu J (2001) In: Ryan MJ (ed.) *Anuran Communication*. Smithsonian Institution Press. Washington and London.

Capítulo 2

- Altmann, J. 1974. Observational Study of Behavior: Sampling Methods. *Behavior*, 49(3/4):227–267.
- Araújo, C.B; Marcondes-Machado, L.O; Vieliard, J.M.E. 2011. Vocal repertoire of the yellow-faced parrot (*Alipiopsitta xanthops*). *The Wilson Journal of Ornithology* 123(3):603–608.
- Charif, R.A; Waack, A.M; Strickman, L.M. 2010. *Raven Pro 1.4 User's Manual*. Cornell Lab of Ornithology, Ithaca, NY.
- Caramaschi, U & Feio, R.N. 1990. A new species of *Hyla* (Anura, Hylidae) from southern Minas Gerais, Brazil. *Copeia* 1990(2): 542–546.
- Carvalho Jr., R.R; Galindo, C.A.B; Nascimento, L.B. 2006. Notes on the courtship behavior of *Aplastodiscus arildae* (Cruz & Peixoto, 1985) at an urban forest fragment in southeastern Brazil (Amphibia, Anura, Hylidae). *Arquivos do Museu Nacional, Rio de Janeiro*. 64(3): 247–254.

- Carvalho, T.R; Giaretta, A.A; Magrini, L. 2012. A new species of the *Bokermannohyla circumdata* group (Anura: Hylidae) from southeastern Brazil, with bioacoustic data on even species of the genus. *Zootaxa* 3321: 37–55.
- Centeno, F.C; Pinheiro, P.D; Andrade, D.V. 2015. Courtship behavior of *Bokermannohyla alvarengai*, a waltzing anuran. *Herpetological Review* 46(2): 166–168.
- Duellman, W.E. & Trueb, L. 1996. *Biology of Amphibians*. The Johns Hopkins University Press. Baltimore and London. 670 p.
- Duellman, W.E; Marrion, A.B; Hedges, S.B. 2016. Phylogenetics, classification, and biogeography of the treefrogs (Amphibia: Anura: Arboranae). *Zootaxa* 4104(1): 001–109.
- Eterovik, P.C & Brandão, R.A. 2001. A description of the tadpoles and advertisement calls of members of the *Hyla pseudopseudis* group. *Journal of Herpetology* 35(3): 442–450.
- Faivovich, J; Haddad, C.F.B; Garcia, P.C.A; Frost, D.R; Campbell, J.A; Wheeler, W.C. 2005. Systematic review of the frog family Hylidae, with special reference to Hylineae: a phylogenetic analysis and taxonomic revision. *Bulletin of the American Museum of Natural History* 294:1–240.
- Frost, D.R. 2016. *Amphibian species of the world: an Online Reference*. Version 6.0 (access: march 28 2016). Electronic database accessible at <http://research.amnh.org/vz/herpetology/amphibia/index.html>. American Museum of Natural History, New York, USA.
- Gaiga, R; Loiola, C; Mângia, S; Pirani, R.M. 2013. Advertisement call and tadpoles of *Bokermannohyla vulcaniae* (Vasconcelos and Giaretta, 2003) (Amphibia: Anura: Hylidae). *South American Journal of Herpetology* 8(2): 127–131.
- Haddad, C.F.B; Faivovich, J; Garcia, P.C.A. 2005. The specialized reproductive mode of the treefrog *Aplastodiscus perviridis* (Anura: Hylidae). *Amphibia–Reptilia* 26: 87–92.
- Haddad, C.F.B. & Prado, C.P.A. 2005. Reproductive modes in frogs and their unexpected diversity in the Atlantic forest of Brazil. *BioScience* 55(3): 207–217.
- Haddad, C.F.B; Sawaya, R. 2000. Reproductive modes of Atlantic Forest Hylids frogs: A general overview and the description of a new mode. *Biotropica* 32(4): 862–871.
- Haddad, C.F.B; Toledo, L.F; Prado, C.P.A; Loebmann, D; Gasparini, J.L; Sazima, I. 2013. *Guia dos Anfíbios da Mata Atlântica: Diversidade e Biologia*. Anolis Books. 543 pp.
- Kluge, A.G. 1981. The life history, social organization, and parental behavior of *Hyla rosenbergi* Boulenger, a nest-building gladiator frog. *Miscellaneous Publications Museum of Zoology, University of Michigan* 160: 1–170.


- Lima, N.G.S; Gontijo, A.S.B; Eterovick, P.C. 2014. Breeding behavior of *Bokermannohyla nanuzae* (Anura: Hylidae) at an Atlantic Forest site in southeastern Brazil. *Journal of Natural History* 48(23-24): 1439–1452.
- Nali, R.C. & Prado, C.P.A. 2012. Habitat use, reproductive traits and social interactions in a stream-dweller treefrog endemic to the Brazilian Cerrado. *Amphibia-Reptilia* 33: 337–347.
- Napoli, M.F & Caramaschi, U. 2004. Two new species of the *Hyla circumdata* group from Serra do Mar and Serra da Mantiqueira, Southeastern Brazil, with description of the advertisement call of *Hyla ibitipoca* (Anura, Hylidae). *Copeia* 2004(3): 534–545.
- Napoli, M.F & Pimenta, B.V.S. 2003. Nova espécie do grupo *Hyla circumdata* (Cope, 1870) do sul da Bahia, Brasil (Amphibia, Anura, Hylidae). *Arquivos do Museu Nacional* 61(3): 189–194.
- Napoli, M.F & Pimenta, B.V.S. 2009. A new species of the *Bokermannohyla circumdata* group (Anura: Hylidae) from the coastal forest of Bahia, Northeastern Brazil. *Copeia* 2009(4):674–683.
- Napoli, M.F & Juncá, F.A. 2006. A new species of the *Bokermannohyla circumdata* group (Amphibia: Anura: Hylidae) from Chapada Diamantina, State of Bahia, Brazil. *Zootaxa* 1244: 57–68.
- Pombal Jr., J.P & Haddad, CFB. 1993. *Hyla luctuosa*, a New Treefrog from Southeastern Brazil (Amphibia: Hylidae). *Herpetologica* 49(1):16–21.
- Robillard, T; Höbel, G; Gerhardt, H.C. 2006. Evolution of the advertisement signals in North American hylid frogs: vocalizations as end-products of calling behavior. *Cladistics* 22:533–545.
- Toledo, L.F; Martins, I.A; Bruschi, D.P; Passos, M.A; Alexandre, C; Haddad, C.F.B. 2015. The anuran calling repertoire in the light of social context. *Acta Ethol.* 18:87–99.
- Vitt, L.J & Caldwell, J.P. 2009. *Herpetology: An introductory biology of amphibians and reptiles*. Third Edition. Elsevier Inc. 697 pp.
- Wells, K.D. 1977. The social behaviour of anuran amphibians. *Animal Behavior* 25: 666–693.
- Wells, K.D. 2007. *The Ecology and Behavior of amphibians*. The University of Chicago Press. Chicago and London. 1148 pp.
- Zina, J. & Haddad, C.F.B. 2007. Courtship behavior of two treefrog species, *Aplastodiscus arildae* and *A. leucopygius* (Anura: Hylidae), from the Atlantic Forest, southeastern Brazil. *Herpetological Review* 38(3): 282–285.


ANEXOS

Profa. Dra. Rachel Meneguello
Presidente
Comissão Central de Pós-Graduação
Declaração

As cópias de artigos de minha autoria ou de minha co-autoria, já publicados ou submetidos para publicação em revistas científicas ou anais de congressos sujeitos a arbitragem, que constam da minha Dissertação/Tese de Mestrado/Doutorado, intitulada **"Comunicação acústica em anuros com especial enfoque para: evolução de perda da vocalização de anúncio e presença de vocalização de corte em *Bokermannohyla luctuosa*"**, não infringem os dispositivos da Lei n.º 9.610/98, nem o direito autoral de qualquer editora.

Campinas, 08/06/2016

Assinatura : 
Nome do(a) autor(a): **Camila Inés Zornosa Torres**
RG n.º V996846-U

Assinatura : 
Nome do(a) orientador(a): **Luís Felipe Toledo**
RG n.º 28.465.361-5



COORDENADORIA DE PÓS-GRADUAÇÃO
INSTITUTO DE BIOLOGIA
Universidade Estadual de Campinas
Caixa Postal 6109. 13083-970, Campinas, SP, Brasil
Fone (19) 3521-6378. email: cpgib@unicamp.br



DECLARAÇÃO

Em observância ao §5º do Artigo 1º da Informação CCPG-UNICAMP/001/15, referente a Bioética e Biossegurança, declaro que o conteúdo de minha Dissertação de Mestrado, intitulada "*Comunicação acústica em anuros com especial enfoque para: evolução de perda da vocalização de anúncio e presença de vocalização de corte em Bokermannohyla luctuosa*", desenvolvida no Programa de Pós-Graduação em Ecologia do Instituto de Biologia da Unicamp, não versa sobre pesquisa envolvendo seres humanos, animais ou temas afetos a Biossegurança.

Assinatura: _____

Nome do(a) aluno(a): Camila Inés Zornosa Torres

Assinatura: _____

Nome do(a) orientador(a): Luís Felipe Toledo

Data: 08/06/2016